

A new contribution to the nuclear modification factor of non-photonic electrons in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV

G. Martínez-García^{1*}, S. Gadrat^{1†} and P. Crochet^{2‡}

¹*SUBATECH (IN2P3/CNRS - Ecole des Mines - Université de Nantes) Nantes, France*

²*Laboratoire de Physique Corpusculaire (IN2P3/CNRS - Université Blaise Pascal)*

Clermont-Ferrand, France

(February 2, 2008)

Abstract

We investigate the effect of the so-called anomalous baryon/meson enhancement to the nuclear modification factor of non-photonic electrons in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV. It is demonstrated that an enhancement of the charm baryon/meson ratio, as it is observed for non-strange and strange hadrons, can be responsible for part of the amplitude of the nuclear modification factor of non-photonic electrons. About half of the measured suppression of non-photonic electrons in the $2 - 4$ GeV/c p_t range can be explained by a charm baryon/meson enhancement of 5. This contribution to the non-photonic electron nuclear modification factor has nothing to do with heavy quark energy loss.

*Gines.Martinez@subatech.in2p3.fr

†Sebastien.Gadrat@subatech.in2p3.fr

‡Philippe.Crochet@clermont.in2p3.fr

One of the most robust experimental evidence for the creation of a new state of matter in heavy-ion collisions at the Relativistic Heavy Ion Collider (RHIC) is the large suppression of light hadrons at high transverse momentum (p_t) [1]. This phenomenon is well reproduced by models which take into account the radiative energy loss of high p_t light quarks and gluons propagating through a dense medium of colored quarks and gluons [2]. Further insights into the underlying mechanism can be obtained from the study of heavy hadrons. Indeed, heavy hadrons originate from the fragmentation of (heavy) quarks in contrast to light hadrons which are predominantly produced by gluons. Quarks are supposed to lose less energy than gluons in the medium due to a smaller color charge coupling. In addition, radiative energy loss was predicted to be substantially smaller for heavy quarks as compared to light quarks because of the so called “dead-cone” effect which limits the medium induced radiative energy loss at forward angles [3]. Surprisingly, recent data from the PHENIX and the STAR collaborations in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV show that the quenching of non-photonic electrons lies beyond theoretical expectations [4–6] and is as large as that of light mesons. This is difficult to understand under the common assumption that non-photonic electrons originate predominantly from the decay of D and B mesons, the latter originating themselves from heavy (c and b) quarks hadronization. Reconciling these data with model predictions is a real challenge which triggers a lot of theoretical activities nowadays. Only models which assume a very large medium opacity [7] or an additional collisional energy loss [8,9] or a very early fragmentation of heavy quarks in the medium [10] can describe successfully the data (for a recent review, see [11]). It is worth noticing that, in contrast to light hadrons, the heavy flavor quenching is, so far, not measured experimentally through identified particles, but in an inclusive way via the nuclear modification factor (R_{AA}) of non-photonic electrons¹.

In this paper, we point out the possibility that part of the explanation for the strong

¹The nuclear modification factor of non-photonic electrons is obtained from the p_t distributions

suppression of non-photonic electrons might be due to another source of electrons namely charmed baryons. Indeed, whereas light mesons are largely suppressed in heavy ion collisions at RHIC, the suppression of non-strange and strange baryons is observed to be much less in the intermediate p_t range ($2 < p_t < 4$ GeV/c) [12]. This is commonly referred to as the anomalous baryon/meson enhancement. It could challenge our understanding of jet quenching at a first glance but brings however another piece of evidence for the formation of a deconfined medium. This anomalous baryon/meson enhancement is well understood in the framework of the recombination model which assumes that, at moderate p_t , hadronization occurs via the coalescence of “free” quarks (and anti-quarks) [13–18]. In the following, we show that an anomalous baryon/meson enhancement for charm hadrons leads naturally to a non-photonic electron R_{AA} smaller than one. This is essentially due to a smaller semi-leptonic decay branching ratio and to a softer decay lepton spectrum of charm baryons as compared to charm mesons. As a consequence, a fraction of the deviation from the experimentally measured R_{AA} of non-photonic electrons should not be attributed to energy loss. After the description of the physical process invoked here, we present simple PYTHIA-based simulations to illustrate the magnitude of the effect in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV and we compare our results to the PHENIX data.

The main assumption we put forward is that, in a deconfined medium, charm baryon production is enhanced relative to charm meson production, as compared to the vacuum. This assumption is qualitatively justified in the framework of the recombination model. Although this model does not provide predictions on charm hadron production yet, it successfully describes the baryon/meson enhancement measured in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV.

of non-photonic electrons in AA collisions (dN_{AA}^e/dp_t) and in pp collisions (dN_{pp}^e/dp_t) as:

$$R_{AA} = \frac{dN_{AA}^e/dp_t}{\langle N_{coll}^{AA} \rangle dN_{pp}^e/dp_t}$$

where $\langle N_{coll}^{AA} \rangle$ is the average number of nucleon-nucleon collisions corresponding to a given centrality class.

A relatively good agreement is obtained not only for the light hadron ratio p/π^+ , but also for heavier hadron ratios such as Λ/K_s^0 and Ω/ϕ [21]. However, extrapolating these results to charm hadrons is not straightforward because the mass of the charm quark is much larger than that of light and strange quarks. The consequences are threefold as far as the recombination mechanism is concerned. First, whereas the p_t of a light baryon (meson) amounts to 3 (2) times the initial p_t of its valence quarks, the p_t of a (single) charm baryon or a charm meson is likely to be very close to that of the charm quark. Secondly, considering a light quark and a heavy quark with the same velocity (which is the essential requirement for the coalescence process to take place [22]), the heavy quark momentum is much larger than that of light partons. As a consequence, one can expect the enhancement of charm baryon/meson to appear at higher p_t . The recombination model indeed predicts, for non-strange and strange hadrons, that the heavier the hadron, the larger the p_t of the baryon/meson enhancement [21]. Finally, the fragmentation time of heavy quarks is small as compared to light quarks. According to [10], the formation time of a 10 GeV/c pion, D meson and B meson is 20, 1.5 and 0.4 fm/c, respectively and it is as small as ~ 3 fm/c for a Λ_c with $p_t = 20 - 30$ GeV/c. Due to these considerations, it is obvious that the baryon/meson enhancement for non-charm hadrons and charm hadrons can be significantly different. We stress that our aim is neither to investigate these differences nor to quantify the magnitude of the charm baryon/meson enhancement. This would require a sophisticated theoretical investigation which goes clearly beyond the scope of this paper and, to our knowledge, no information on this topic is available in the literature yet. Therefore, in the following, we only assume that, in view of experimental results on the baryon/meson enhancement for non-strange and strange hadrons, a similar enhancement is a priori conceivable for charm hadrons. Remarkably, such an enhancement has strong implications on the nuclear modification factor of non-photonic electrons. It leads to a decrease of the yield of non-photonic electrons in AA collisions because, as shown in Tab. I, the inclusive semi-leptonic decay branching ratio of charm baryons is smaller than that of charm mesons (and the hadronization process does not change the total charm quark number). Therefore, the

nuclear modification factor of non-photonic electrons should decrease as well. This can be easily illustrated in the following way. Assuming that a AA collision is a perfect sum of pp collisions (i.e. $R_{AA} = 1$) and that the relative yields of D mesons are the same in pp and in AA collisions, a p_t integrated R_{AA} can be calculated for different Λ_c/D enhancement factors $\mathcal{C} = N_{\Lambda_c, \bar{\Lambda}_c}/N_D = (N_{\Lambda_c} + N_{\bar{\Lambda}_c})/(N_{D^+} + N_{D^-} + N_{D^0} + N_{\bar{D}^0} + N_{D_s^+} + N_{D_s^-})$ according to

$$R_{AA} = \frac{1 + N_{\Lambda_c, \bar{\Lambda}_c}/N_D}{1 + \mathcal{C}(N_{\Lambda_c, \bar{\Lambda}_c}/N_D)} \frac{1 + \mathcal{C}(N_{\Lambda_c \rightarrow e}/N_{D \rightarrow e})}{1 + N_{\Lambda_c \rightarrow e}/N_{D \rightarrow e}} \quad (1)$$

where

$$N_{\Lambda_c \rightarrow e}/N_{D \rightarrow e} = \frac{(N_{\Lambda_c, \bar{\Lambda}_c}/N_D)BR_{\Lambda_c, \bar{\Lambda}_c}}{(N_{D^\pm}/N_D)BR_{D^\pm} + (N_{D^0, \bar{D}^0}/N_D)BR_{D^0, \bar{D}^0} + (N_{D_s^\pm}/N_D)BR_{D_s^\pm}}. \quad (2)$$

N is the charm hadron yield and BR is the hadron semi-leptonic decay branching ratio. According to Tab. I, $N_{\Lambda_c, \bar{\Lambda}_c}/N_D = 7.3\%$, $N_{D^\pm}/N_D = 21\%$, $N_{D^0, \bar{D}^0}/N_D = 67\%$, and $N_{D_s^\pm}/N_D = 12\%$ such that $N_{\Lambda_c \rightarrow e}/N_{D \rightarrow e} = 3.63\%$. Enhancement factors $\mathcal{C}$ of 5 and 12 therefore lead to a R_{AA} of 0.90 and 0.79 respectively. It is shown below that the effect is further amplified in the intermediate p_t range as a result of a softer decay electron spectrum of Λ_c as compared to that of D mesons.

In order to investigate this effect in more details, we have performed simple simulations with the PYTHIA-6.152 [23] event generator. The PYTHIA input parameters were first tuned according to [20] and the PHENIX acceptance cut ($|\eta| < 0.35$) was applied in order to correctly reproduce the p_t distribution of non-photonic electrons measured in pp collisions at $\sqrt{s} = 200$ GeV [24]. As it can be seen from Fig. 1, the agreement between the simulation and the data is rather good except at very low p_t and in the high p_t region where the simulation under-predicts the data. The disagreement in the high p_t region is due to the fact that we consider, in the simulation, only electrons from charm decay and neglect electrons from bottom decay which are expected to dominate the spectrum for $p_t \gtrsim 4$ GeV/c [24]. On the other hand, it can be observed from Tab. I that the Λ_c/D ratio amounts to 7.3% (in 4π) and, as mentionned above, to 3.63% after convolution of the species yields with their corresponding semi-leptonic decay branching ratio. Figure 2 shows that, in the $2 - 4$ GeV/c

p_t region of interest discussed hereafter, this ratio is even smaller ($\sim 1.5\%$) because the decay electron spectrum of Λ_c is softer than that of D mesons. In this respect, an enhancement of the Λ_c/D ratio in central heavy ion collisions induces two different mechanisms of non-photonic electron yield suppression: i) integrated yields are suppressed due to the lower semi-electronic branching ratio of the Λ_c and ii) the softer electron p_t distribution from Λ_c decay leads to an additional suppression at intermediate p_t .

We then re-evaluate the non-photonic electron decay spectrum after introducing a Λ_c/D enhancement chosen to 5 and 12. This enhancement is assumed to be flat in p_t and it is applied such that the p_t -differential charm cross-section is conserved. The latter is an arbitrary choice that could be justified since most of the charm hadron transverse momentum is given by the charm quark whatever, baryon or meson, this hadron is. We finally compute the R_{AA} ratio from the previous non-photonic electron p_t spectra neglecting all effects like shadowing or quenching but assuming that the only medium induced effect is the Λ_c/D enhancement. The nuclear modification factor is therefore simply obtained from the ratio of the non-photonic electron p_t distribution with a Λ_c/D enhancement to the original distribution. By doing so, one implicitly takes into account the p_t dependence of the Λ_c/D ratio which is not flat (Fig. 2). The results are shown in Fig. 3 together with the PHENIX data. The simulated R_{AA} ratio is shown only for $2 < p_t < 4$ GeV/c since shadowing and electrons from bottom decay, which play an important role at lower and higher p_t respectively, are not considered in the simulation. One can see from Fig. 3 that a Λ_c/D enhancement of 5 can already explain $\sim 50\%$ of the suppression of non-photonic electrons in the $2 - 4$ GeV/c p_t range. For illustration, we show that, in the extreme scenario of an enhancement of 12, the data can be satisfactorily described without invoking any heavy-quark energy loss. The shape of the simulated R_{AA} is obviously determined by the p_t distribution of Λ_c/D enhancement that we have assumed to be flat. Note that the values of R_{AA} at $p_t \sim 3$ GeV/c can be roughly obtained as well from Fig. 2 and Eq. 1 and 2, assuming that the decay electrons at $p_t \sim 3$ GeV/c originate from charm hadrons with $p_t \gtrsim 4$ GeV/c.

In summary, we have shown that an enhancement of the Λ_c/D ratio in heavy ion colli-

sions has dramatic consequences on the nuclear modification factor of non-photonic electrons. Such an enhancement, which is observed for non-strange and strange hadrons, would significantly lower the R_{AA} of non-photonic electrons (or muons) at intermediate p_t . This results from i) the semi-leptonic decay branching ratio of charm baryons which is smaller than that of charm mesons and ii) the decay lepton spectrum from charm baryons which is softer than that of charm mesons. We conclude that it is therefore premature to interpret the non-photonic electron R_{AA} data before a possible enhancement of the Λ_c/D ratio is measured experimentally and/or investigated theoretically. Heavy quark energy loss can be studied in a much cleaner way from the nuclear modification factor of exclusively reconstructed charm hadrons. Such measurements should be possible with the RHIC-II experiments [25] and with the ALICE experiment at the LHC [26]. We finally note that the Λ_c/D enhancement can possibly influence the elliptic flow of non-photonic electrons as well.

NOTE

After the publication of the present work in arXiv we have realized that a similar work has already been published by P. Sorensen and X. Dong in PRC 74 (2006) 024902. However, the assumptions made in the two approaches are different and lead to different conclusions. In PRC 74 (2006) 024902, assuming a Λ_c/D ratio of 1.7, similar to the measured Λ/K_s^0 ratio, about 35% of the of non-photonic electron R_{AA} data is reproduced. In our approach, the R_{AA} data are reproduced by using the Λ_c and D spectra from PYTHIA and assuming a Λ_c/D ratio of ~ 1 (i.e 12×0.08).

ACKNOWLEDGMENTS

We gratefully acknowledge Anton Andronic, Pol-Bernard Gossiaux and Stéphane Peigné for carefully reading the manuscript and for valuable suggestions.

REFERENCES

- [1] see e.g. J. Adams *et al.* [STAR Collaboration], Phys. Rev. Lett. **91**, 172302 (2003) [arXiv:nucl-ex/0305015], C. Adler *et al.* [STAR Collaboration], Phys. Rev. Lett. **89**, 202301 (2002) [arXiv:nucl-ex/0206011], S. S. Adler [PHENIX Collaboration], arXiv:nucl-ex/0611006, S. S. Adler [PHENIX Collaboration], arXiv:nucl-ex/0611007, I. Arsene [BRAHMS Collaboration], arXiv:nucl-ex/0610021, B. Alver *et al.* [PHOBOS Collaboration], Phys. Rev. Lett. **96**, 212301 (2006) [arXiv:nucl-ex/0512016].
- [2] For reviews, see e.g. C. A. Salgado, Nucl. Phys. A **774**, 267 (2006) [arXiv:hep-ph/0510062]; X. N. Wang, Nucl. Phys. A **774**, 215 (2006) [arXiv:nucl-th/0511001] and references therein.
- [3] Y. L. Dokshitzer and D. E. Kharzeev, Phys. Lett. B **519**, 199 (2001) [arXiv:hep-ph/0106202].
- [4] A. Adare *et al.* [PHENIX Collaboration], arXiv:nucl-ex/0611018.
- [5] S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **96**, 032301 (2006) [arXiv:nucl-ex/0510047].
- [6] B. I. Abelev *et al.* [STAR Collaboration], arXiv:nucl-ex/0607012.
- [7] N. Armesto, M. Cacciari, A. Dainese, C. A. Salgado and U. A. Wiedemann, Phys. Lett. B **637**, 362 (2006) [arXiv:hep-ph/0511257].
- [8] M. Djordjevic, J. Phys. G **32**, S333 (2006) [arXiv:nucl-th/0610054].
- [9] H. van Hees, V. Greco and R. Rapp, Phys. Rev. C **73**, 034913 (2006) [arXiv:nucl-th/0508055].
- [10] A. Adil and I. Vitev, arXiv:hep-ph/0611109.
- [11] R. Auerbeck, proceedings of Quark Matter 2006, 19th Conference on Ultra-Relativistic Nucleus-Nucleus Collisions, Shanghai, November 14-20, 2006, to appear in J. Phys. G.

- [12] see e.g. B. I. Abelev *et al.* [STAR Collaboration], Phys. Rev. Lett. **97**, 152301 (2006) [arXiv:nucl-ex/0606003], J. Adams *et al.* [STAR Collaborations], arXiv:nucl-ex/0601042 S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **91**, 172301 (2003) [arXiv:nucl-ex/0305036], S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. C **69**, 034909 (2004) [arXiv:nucl-ex/0307022], I. Arsene *et al.* [BRAHMS Collaboration], Phys. Rev. C **72**, 014908 (2005) [arXiv:nucl-ex/0503010].
- [13] R. J. Fries, B. Muller, C. Nonaka and S. A. Bass, Phys. Rev. Lett. **90**, 202303 (2003) [arXiv:nucl-th/0301087].
- [14] R. J. Fries, B. Muller, C. Nonaka and S. A. Bass, Phys. Rev. C **68**, 044902 (2003) [arXiv:nucl-th/0306027].
- [15] V. Greco, C. M. Ko and P. Levai, Phys. Rev. Lett. **90**, 202302 (2003) [arXiv:nucl-th/0301093].
- [16] V. Greco, C. M. Ko and P. Levai, Phys. Rev. C **68**, 034904 (2003) [arXiv:nucl-th/0305024].
- [17] R. C. Hwa and C. B. Yang, Phys. Rev. C **70**, 024904 (2004) [arXiv:hep-ph/0312271].
- [18] R. C. Hwa and C. B. Yang, Phys. Rev. C **67**, 034902 (2003) [arXiv:nucl-th/0211010].
- [19] W. M. Yao *et al.* [Particle Data Group], J. Phys. G **33**, 1 (2006).
- [20] K. Adcox *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **88**, 192303 (2002) [arXiv:nucl-ex/0202002].
- [21] S. Blyth for the STAR Collaboration, proceedings of Quark Matter 2006, 19th Conference on Ultra-Relativistic Nucleus-Nucleus Collisions, Shanghai, November 14-20, 2006, to appear in J. Phys. G, [arXiv:nucl-ex/0701052].
- [22] Z. W. Lin and D. Molnar, Phys. Rev. C **68**, 044901 (2003) [arXiv:nucl-th/0304045].
- [23] T. Sjöstrand, Comput. Phys. Commun. **82**, 74 (1994).

- [24] A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **97**, 252002 (2006) [arXiv:hep-ex/0609010].
- [25] B. Jacak, proceedings of Quark Matter 2006, 19th Conference on Ultra-Relativistic Nucleus-Nucleus Collisions, Shanghai, November 14-20, 2006, to appear in J. Phys. G.
- [26] B. Alessandro *et al.* [ALICE Collaboration], J. Phys. G **32**, 1295 (2006).

TABLES

TABLE I. Inclusive decay branching ratio (BR) of charm hadrons into $e + \text{anything}$ [19] and yield (N) of charm hadrons (in 4π) in pp collisions at $\sqrt{s} = 200$ GeV from the present PYTHIA simulation using input parameters as described in [20]. The total cross-section for charm production is normalized to the experimental value obtained in [20]. N_{Λ_c} and $N_{\bar{\Lambda}_c}$ include primarily produced Λ_c and $\bar{\Lambda}_c$ as well as those from Σ_c and $\bar{\Sigma}_c$ decay.

Hadron	D^+	D^-	D^0	$\bar{D}^0$	D_s^+	D_s^-	Λ_c	$\bar{\Lambda}_c$
BR (%)	17.2 ± 1.9		6.71 ± 0.29		8_{-5}^{+6}		4.5 ± 1.7	
N ($\times 10^{-3}$)	3.00	3.07	9.31	9.85	1.82	1.60	1.23	0.85

FIGURES




FIG. 1. Invariant differential cross-section of non-photonics electrons (dots) measured in pp collisions at $\sqrt{s} = 200$ GeV [24]. The solid curve shows the result of the PYTHIA simulation as described in the text. The simulated spectrum is normalized from the integration of the measured spectrum in the range $1.4 < p_t < 4$ GeV/c.




FIG. 2. Charm baryons over charm mesons (dots) and decay electrons from charm baryons over decay electrons from charm mesons (squares) versus p_t , from the PYTHIA simulation described in the text.




FIG. 3. Nuclear modification factor of non-photonic electrons (dots) measured in central (0-10%) Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV [4]. The solid and dotted curves correspond to the results of the simulation described in the text for a Λ_c/D enhancement factor of 5 and 12 respectively.